

RESEARCH PAPER

Xylem vulnerability to cavitation can be accurately characterised in species with long vessels using a centrifuge method

M. F. Tobin¹, R. B. Pratt¹, A. L. Jacobsen¹ & M. E. De Guzman²

¹ Department of Biology, California State University, Bakersfield, Bakersfield, CA, USA

² Department of Botany and Plant Sciences, University of California, Riverside, Riverside, CA, USA

Keywords

Cavitation resistance; centrifuge; dehydration; embolism; vessel length; vulnerability curves; xylem.

Correspondence

M. Tobin, Department of Biology, California State University, Bakersfield, 9001 Stockdale Highway, Bakersfield, CA 93311, USA.

E-mail: miketobin@alumni.reed.edu

Editor

J. Sparks

Received: 6 May 2012; Accepted: 23 August 2012

doi:10.1111/j.1438-8677.2012.00678.x

ABSTRACT

Vulnerability to cavitation curves describe the decrease in xylem hydraulic conductivity as xylem pressure declines. Several techniques for constructing vulnerability curves use centrifugal force to induce negative xylem pressure in stem or root segments. Centrifuge vulnerability curves constructed for long-vesselled species have been hypothesised to overestimate xylem vulnerability to cavitation due to increased vulnerability of vessels cut open at stem ends that extend to the middle or throughout segments. We tested two key predictions of this hypothesis: (i) centrifugation induces greater embolism than dehydration in long-vesselled species, and (ii) the proportion of open vessels changes centrifuge vulnerability curves. Centrifuge and dehydration vulnerability curves were compared for a long- and short-vesselled species. The effect of open vessels was tested in four species by comparing centrifuge vulnerability curves for stems of two lengths. Centrifuge and dehydration vulnerability curves agreed well for the long- and short-vesselled species. Centrifuge vulnerability curves constructed using two stem lengths were similar. Also, the distribution of embolism along the length of centrifuged stems matched the theoretical pressure profile induced by centrifugation. We conclude that vulnerability to cavitation can be accurately characterised with vulnerability curves constructed using a centrifuge technique, even in long-vesselled species.

INTRODUCTION

Plants transport water from the soil to leaves along a negative pressure gradient according to the cohesion–tension theory (Dixon & Joly 1895). While energetically efficient, transport of water under negative pressure is susceptible to cavitation, and subsequent embolism formation can block water flow through xylem conduits, thereby reducing hydraulic efficiency (Sperry & Tyree 1988; Tyree & Sperry 1989). Species differ broadly in their vulnerability to cavitation (Pockman & Sperry 2000; Jacobsen *et al.* 2007b), as do plant communities (Jacobsen *et al.* 2007a). Moreover, vulnerability to cavitation has been linked to drought resistance and survival (Pratt *et al.* 2008; Kursar *et al.* 2009), life-history characteristics (Pratt *et al.* 2007) and stomatal behaviour (Brodribb & Holbrook 2004), and it is evolutionarily correlated with many other functional traits (Maherali *et al.* 2004; Jacobsen *et al.* 2007b). The broad significance of the vulnerability of plants to cavitation has made it the focus of intense study.

The vulnerability of plants to cavitation is often characterised by constructing vulnerability curves, which describe the decline in xylem hydraulic conductivity (K_h) due to embolism formation in response to increasingly negative xylem pressure. Declines in K_h are typically expressed relative to the maximum sample K_h as the percentage loss of conductivity (PLC), and comparisons of the vulnerability to cavitation among species are made using the xylem pressure at 50% loss of conductivity (P_{50}). An early method for constructing vulnerability curves

involved collecting large branches and equilibrating them to a range of xylem pressures by repeated cycles of drying and equilibration while measuring PLC of stems on those branches at each equilibration xylem pressure (Sperry *et al.* 1988b). Although labour intensive, this method has the advantage that negative xylem pressure is generated in a similar manner to an intact plant, and it remains a standard to which other methods are compared.


Several techniques for constructing vulnerability curves rely on centrifugal force to induce a range of negative xylem pressures in stem or root segments (Pockman *et al.* 1995; Alder *et al.* 1997; Cochard *et al.* 2005; Li *et al.* 2008). The negative xylem pressure induced by centrifugation depends on segment length and angular velocity. The pressure profile along the segment is most negative at its centre and becomes less negative towards both ends. All centrifuge techniques include these features, but techniques differ in other attributes.

In the technique used in this study, described by Alder *et al.* (1997), segments are mounted into a custom centrifuge rotor configured so that when it is spinning, segment ends are submerged in solution maintained at equal levels. Because solution levels are equal, no flow through the segment occurs during centrifugation. In this technique, segments are removed from the centrifuge to measure K_h under a low positive pressure (Sperry *et al.* 1988a). Through repeatedly centrifuging segments to more negative xylem pressures and measuring K_h between spins, a complete vulnerability curve can be constructed.

1 An alternate centrifuge technique relying on the 'Cavitron'
 2 rotor design (Cochard 2002; Cochard *et al.* 2005) enables the
 3 K_h of a segment to be measured while it is spinning in the cen-
 4 trifuge. This is accomplished by increasing the solution level at
 5 one end of the segment to create a pressure gradient across the
 6 segment that drives flow during centrifugation. This technique
 7 has the advantage of measuring K_h while the xylem is under
 8 negative pressure, as occurs *in vivo*. Although similar in con-
 9 cept to the Cavitron technique, the centrifuge technique
 10 described by Li *et al.* (2008) used a different rotor design.
 11 These distinctions are important, since differences in rotor
 12 design may make some rotors more prone to artefacts than
 13 others (Sperry *et al.* 2012).

14 Two studies have questioned whether vulnerability curves
 15 constructed with a centrifuge method accurately characterise
 16 the vulnerability to cavitation for long-vesselled species (Choat
 17 *et al.* 2010; Cochard *et al.* 2010). Choat *et al.* (2010) found
 18 that centrifuge vulnerability curves constructed for *Vitis vinifera*
 19 (grapevine) using the technique described in Alder *et al.*
 20 (1997) did not agree with vulnerability curves constructed
 21 using other methods. In a second study, Cochard *et al.* (2010)
 22 showed that vulnerability curves for a long-vesselled species
 23 constructed with the Cavitron centrifuge technique differed
 24 markedly from dehydration curves, and were shifted by altering
 25 the length of the stem segment (Fig. 1). The authors of both
 26 studies hypothesised that vessels cut open at segment ends and
 27 extending to the middle or entirely through segments (open
 28 vessels) are more vulnerable to cavitation during centrifugation
 29 than they would be *in situ* during dehydration. If supported,
 30 this open-vessel artefact hypothesis raises doubts about the
 31 interpretation of centrifuge vulnerability curves constructed for
 32 species with many open vessels. However, reinterpretation of
 33 centrifuge vulnerability curves for long-vesselled species may
 34 be unwarranted because of recently reported methodological
 35 challenges associated with *V. vinifera* (Jacobsen & Pratt 2012),
 36 and the substantial differences between the Cavitron rotor
 37 design and measurement protocol and other centrifuge
 38 techniques.

39 We tested key predictions of the open-vessel artefact hypoth-
 40 esis to determine the reliability of vulnerability curves con-
 41 structed with the centrifuge technique described in Alder *et al.*
 42 (1997). To test whether open vessels created during stem
 43



57 **Fig. 1.** The percentage loss of hydraulic conductivity (PLC) as a function of
 58 xylem pressure reported by Cochard *et al.* (2010) for *Quercus robur*, show-
 59 ing the wide divergence of dehydration and Cavitron curves and between
 60 Cavitron curves constructed with 175 and 275-mm stem segments (data
 61 redrawn from Cochard *et al.* 2010).

segment preparation increased segment vulnerability to cavita-
 tion, we compared laboratory dehydration vulnerability curves
 to matched centrifuge curves for a long-vesselled (*Quercus wis-*
lizeni) and a short-vesselled (*Heteromeles arbutifolia*) shrub
 species. The long-vesselled oak was chosen because it has a
 maximum vessel length of 1.45 m (Jacobsen *et al.* 2007b),
 which is comparable to the 1.34 m maximum vessel length of
 the oak (*Quercus robur*) that Cochard *et al.* (2010) found to
 have widely divergent dehydration and Cavitron vulnerability
 curves (Fig. 1). We constructed these curves by first determin-
 ing the PLC of stems sampled from large branches during labo-
 ratory dehydration, and then flushing and centrifuging the
 stems to a xylem pressure similar to that measured for the large
 branch with stem psychrometers. This allowed us to directly
 compare the level of stem embolism induced by air-drying
 large branches or by centrifugation of stem segments. In addi-
 tion, an acoustic method was used to independently monitor
 cavitation in branches during dehydration (Tyree & Sperry
 1989), and native embolism measurements of plants in the field
 were compared to vulnerability curves.

We also tested the prediction of the open-vessel artefact
 hypothesis that a larger proportion of open vessels in a stem
 segment will increase its vulnerability to cavitation during cen-
 trifugation of a centrifuge vulnerability curve. To test this predic-
 tion, we compared centrifuge vulnerability curves constructed
 with stem segments that differed in length and therefore also
 the proportion of open vessels. Comparisons were made for
 four species, including *Q. robur*, the long-vesselled oak tested
 by Cochard *et al.* (2010) and reported to have Cavitron vulner-
 ability curves that differed depending on segment length
 (Fig. 1), and *V. vinifera*, the species found by Choat *et al.*
 (2010) to have anomalous centrifuge vulnerability curves.

As another test of the centrifuge technique used in this
 study, we characterised the distribution of embolism along the
 length of centrifuged stem segments of a long-vesselled species
 to compare it with the theoretical pressure profile induced by
 centrifugation. The distribution of embolism produced by the
 Cavitron (Fig. 5 in Cochard *et al.* 2010) did not match theoret-
 ical predictions that the level of embolism should be highest at
 the centre of the stem; therefore we tested our technique for a
 similar departure from theoretical predictions.

MATERIAL AND METHODS

Study site and plant material

Chaparral shrub plant material was collected from a site in the
 San Gabriel Mountains of southern California (1040 m a.s.l.;
 34°21.613 N, 118°26.125 W). The site has a mediterranean-
 type climate with hot, dry summers and cool, wet winters and
 a mean annual precipitation of 685 mm from 1996 to 2010.
 Branches of long-vesselled *V. vinifera* L. cv. Glenora were col-
 lected from a mature, irrigated plant growing in Bakersfield,
 CA, USA. For *V. vinifera*, all branches were collected from the
 same individual to control for interplant variability and to
 reduce variability in the number of open vessels not due to dif-
 ferences in sample length. Branches of long-vesselled *Q. robur*
 L. cv. *fastigiata* were collected from plants grown in a green-
 house in Bakersfield with daily irrigation for 8 months, follow-
 ing their purchase as ~1-m plants from Lawyer Nursery Inc.,
 MT, USA.

Two evergreen chaparral shrubs were chosen for the main experiments because of their widely divergent vessel lengths (Table 1): the long-vesselled *Q. wislizeni* var. *frutescens* Engelm. (Fagaceae) and the shorter-vesselled *Heteromeles arbutifolia* (Lindl.) M. Roem. (Rosaceae). Two additional long-vesselled species (mean vessel length >0.10 m), *Rhus ovata* S. Watson (Anacardiaceae) and *Fraxinus dipetala* Hook. and Arn. (Oleaceae), and two shorter-vesselled species (mean vessel length <0.06 m), *Ceanothus leucodermis* Greene (Rhamnaceae) and *Ribes malvaceum* Sm. (Grossulariaceae), were sampled for some experiments. Of these, *F. dipetala* and *R. malvaceum* were deciduous and the rest were evergreen.

We collected large branches (1.8–3.0 m in length) of *Q. wislizeni* and *H. arbutifolia* for laboratory dehydration vulnerability curves from January to June of 2011, before new leaves had flushed. Stem cutting was done under water, branches were enclosed in doubled and humidified plastic bags to reduce transpiration, and the cut end was kept under water during transport to the laboratory. Smaller branches for centrifuge vulnerability curves and a repeated centrifugation experiment (see below) were collected similarly during the same time period, except for *V. vinifera* and *Q. robur*, which were sampled in autumn 2011.

Stem hydraulic conductivity, loss of conductivity and centrifugation

We determined K_h from measurements of the net increase in fluid flow rate through a stem after applying a pressure head (Sperry *et al.* 1988a). We selected stem segments that were ca. 6–8 mm in diameter and prepared them for measurements by cutting segments from sampled stems under water and trimming them to the desired length with a razor blade. To make flow rate measurements, stems were inserted into a tubing apparatus that connected the basal end to an elevated fluid reservoir and the other end to a collection reservoir on an analytical balance (CP124S, Sartorius, Goettingen, Germany) connected to a laptop computer (Inspiron 5000e, Dell Computer Corp., Round Rock, TX, USA) that logged net flow over 10-s intervals. The pressure head was maintained below 2 kPa for stems of *V. vinifera* and below 4 kPa for all other stems to

Table 1. Mean vessel length, percentage of vessels that are cut open at the ends of 140 and 271-mm stem segments and extend entirely through segments, and percentage of vessels open to the middle of segments for *Quercus wislizeni*, *Heteromeles arbutifolia* and *Rhus ovata*.

species	mean vessel length (m)	140-mm stem segments		271-mm stem segments	
		open (%)	open to the middle (%)	open (%)	open to the middle (%)
<i>Quercus wislizeni</i>	0.124 (0.010)	10.7 (1.9)	53.3 (4.1)	1.5 (0.5)	21.4 (3.5)
<i>Heteromeles arbutifolia</i>	0.060 (0.004)	1.1 (0.3)	18.9 (2.7)	0.0 (0.0)	2.5 (0.7)
<i>Rhus ovata</i>	0.143 (0.007)	14.1 (1.4)	60.6 (2.3)	2.4 (0.5)	27.7 (2.4)

Values shown are means and SE in parentheses.

avoid refilling open vessels of large diameters. Stems were kept submerged in water during and between measurements. The perfusing solution used in the tubing apparatus was a 20 mM KCl filtered to 0.1 μ m (inline filter; GE Water and Process Technologies, Trevose, PA, USA) and vacuum-degassed for at least 20 min (Melcher *et al.* 2012). Before and after each measurement of pressure-driven flow, we measured background flow with no pressure head. We calculated K_h as the pressure-driven flow corrected for background flows divided by the pressure gradient (pressure head/stem length). Maximum K_h was measured on stems flushed with the same solution used for K_h measurements at 100 kPa for 1 h to refill embolised vessels. To calculate stem area specific conductivity (K_s), K_h was divided by stem sapwood cross-sectional area, which was calculated from caliper measurements of xylem and pith diameter.

We used centrifugation at a stage in each experiment to induce negative xylem pressure in stem segments (Holbrook *et al.* 1995). We mounted stems in a custom rotor (Alder *et al.* 1997) that enabled them to be spun in a centrifuge (Sorvall RC-5C; Thermo Fisher Scientific, Waltham, MA, USA) with stem ends submerged in solution contained in L-shaped reservoirs. The solution level in the reservoirs was kept even at both ends of the stem, so no flow was induced during centrifugation. During some experiments, we added foam pads to the vertical section of reservoirs to keep the solution in contact with stem ends when the rotor was not spinning (see below, Repeated centrifugation tests).

Laboratory dehydration vulnerability curves

Laboratory dehydration vulnerability curves were completed for 11 large branches of *Q. wislizeni* and three of *H. arbutifolia*. Both species had substantial native embolism (PLC >50% for many stems) that did not reverse during overnight rehydration of branches. The high native PLC made it impossible to fully characterise the vulnerability to cavitation of stems. Therefore, we refilled native embolism by re-cutting branches under water in the laboratory and attaching them to a tubing manifold pressurised to ~15 kPa for about 18 h with the same solution used for K_h measurements. We confirmed removal of native embolism by measuring the PLC of at least two stems per branch.

After refilling, 1–3 stem psychrometers (Stem Hygrometer; PWS Instruments Inc., Guelph, ON, Canada) connected to a data logger (CR7; Campbell Scientific, Logan, UT, USA) were installed on each branch to track branch xylem pressure. The bark was sanded away to expose a small patch of xylem where psychrometers were attached, and silicone grease was used to create an airtight seal. An ultrasonic acoustic emissions transducer (Model I15I; Physical Acoustics Corp., Princeton, NJ, USA) was attached to one of the branches to monitor cavitation as the branch dried. Bark was sanded off the branch where it was attached, and the xylem was covered with silicone grease to facilitate contact with the transducer. A spring-loaded attachment held the transducer against the stem with 30 N of force (Jackson & Grace 1996). The transducer was connected to hardware that conditioned the transducer output so that individual acoustic events were sampled only once, and that amplified the signal by 75 dB. A data logger (21 \times ; Campbell Scientific) sampled this output every second and recorded 5-min sums of acoustic emissions.

1 Branches were equilibrated at increasingly negative xylem
 2 pressure through repeated cycles of drying and equilibration.
 3 Branches were dried by uncovering them for a short period,
 4 then equilibrated by sealing them in double plastic bags, which
 5 were externally covered with moist cotton sheets and more
 6 plastic bags, and allowing them to equilibrate for 16–23 h. Fol-
 7 lowing equilibration, 1–3 stems were removed from each large
 8 branch for PLC measurements, with cuts made under water. At
 9 the same time, we collected up to five branchlets for water
 10 potential measurements with a pressure chamber (PMS Instru-
 11 ment Co., Albany, OR, USA) to compare with stem psychrom-
 12 eter values. We repeated these steps until the stem loss of
 13 hydraulic conductivity was near 100%.

15 Comparison of dehydration and centrifuge methods

17 We constructed centrifuge vulnerability curves using stems
 18 sampled during laboratory dehydration vulnerability curves for
 19 both *Q. wislizeni* and *H. arbutifolia* to test if negative xylem
 20 pressure induced by either air-drying large branches or centri-
 21 fugation of stem segments resulted in different curves. After
 22 the PLC of stems sampled from large branches during a labora-
 23 tory dehydration vulnerability curve was determined, we cen-
 24 trifuged the flushed stems for 5 min to a xylem pressure
 25 similar to that measured for the large branch with psychrome-
 26 ters. The centrifugation-induced PLC of these stems was then
 27 measured, which resulted in a matched pair of centrifuge and
 28 dehydration PLC values at similar water potentials for each
 29 stem segment.

31 Comparison of centrifuge vulnerability curves 32 of two stem lengths

34 We constructed centrifuge vulnerability curves for stems of 140
 35 and 271-mm length for *Q. wislizeni*, *H. arbutifolia*, *Q. robur*
 36 and *V. vinifera* to test the effect of changing the proportion of
 37 open vessels. After an initial flush, we measured the K_h of stems
 38 and then centrifuged them for 5 min ($n = 10$ for 140-mm
 39 stems and $n = 12$ for 271-mm stems of *V. vinifera*, $n = 6$ for
 40 all other species). Stem K_h measurement and centrifugation
 41 was repeated, inducing a more negative xylem pressure with
 42 each centrifugation, and continued until stem loss of hydraulic
 43 conductivity was near 100%.

45 Distribution of centrifugation-induced embolism

47 We measured the distribution of embolism along the length of
 48 stems to compare it with the theoretical negative pressure pro-
 49 file generated during centrifugation. We centrifuged 271-mm
 50 long flushed stems ($n = 3$) of the long-vesselled *R. ovata* at an
 51 angular velocity that induced -2 MPa of pressure at the rota-
 52 tional centre of the stem and reduced entire stem conductivity
 53 by 73–81%. We excised the basal, central and distal 40-mm
 54 segments of these stems and measured their PLC.

56 Repeated centrifugation tests

58 We repeatedly centrifuged stems of all six study species at the
 59 same angular velocity to test for increases in PLC unrelated to
 60 inducing more negative xylem pressure. After an initial flush
 61 and K_h measurement, stems were either centrifuged at

–0.5 MPa for 5 min four times, with K_h measured after each
 centrifugation, or stems were centrifuged once at –0.5 MPa
 for 20 min, with K_h measured afterwards. All stems were then
 flushed, followed by a final K_h measurement. There were three
 stems per treatment, except *R. ovata*, for which there were
 six.

As a follow up, we tested if several treatments reduced the
 increase in PLC observed with repeated centrifugation at the
 same angular velocity. Because the centrifuge reservoir design
 relies on centrifugal force to maintain solution in contact with
 stem ends, they are exposed to air when the rotor is not spin-
 ning. We added foam pads (CoverGirl Make-Up Masters
 Sponge Wedges; Procter & Gamble, Hunt Valley, MD, USA)
 thick enough to be in contact with stem ends to the vertical
 section of reservoirs. The pads were saturated with solution
 and thereby kept the solution in contact with stem ends when
 the rotor was stationary. We tested the effect of foam pads by
 comparing the PLC increase with and without pads for repeat-
 edly centrifuged stems of *R. ovata* ($n = 12$). We also tested the
 addition of 10 min at 0.1 MPa to the end of centrifugation as a
 ‘relaxation’ period on *Q. wislizeni* stems ($n = 3$). Finally,
 recent concerns about solution gas content (Espino & Schenk
 2011) led us to test if additional degassing of the perfusing
 solution reduced increases in PLC with repeated centrifugation
 for *R. ovata* stems ($n = 3$). After our standard vacuum degas-
 sing of 10 l of continuously stirred solution for at least 20 min
 at an absolute pressure of ~10 kPa generated with a vacuum
 pump (model 5KH33DN16X; Millipore, GE, Billerica, MA,
 USA), we further vacuum degassed the solution using a mem-
 brane contactor (Liqui-Cel mini-module 1.7 × 5.5; Membrana,
 Charlotte, NC, USA).

Native PLC

We measured native PLC in *Q. wislizeni* at the end of the
 summer dry period in 2009, and again during the rainy season
 in early spring of 2010. For summer measurements, we col-
 lected large branches ($n = 6$) by cutting them in air near mid-
 day, and brought them to the laboratory sealed in doubled
 and humidified plastic bags. After an equilibration period, we
 determined branch water potential from branchlets using the
 pressure chamber and measured PLC of stems cut under
 water from the large branches at a distance from the original
 cut end that exceeded maximum vessel length. Spring mea-
 surements of native PLC were done in the field on stems col-
 lected pre-dawn, with cuts made under water. Pre-dawn water
 potential for each sampled shrub ($n = 6$) was determined
 from branchlets collected with stems. Laboratory and field-
 portable conductivity systems differed primarily in the
 method used to quantify flow, which was done in the field by
 timing the rate at which solution volume increased in a
 0.1 ml pipette.

Vessel length distributions

Vessel length distribution was measured on six stem segments
 per species for the six chaparral study species using a silicon
 injection method (Sperry *et al.* 2005; Wheeler *et al.* 2005).
 We used stem segments that were 6–8 mm in diameter and
 longer than 250 mm. Prior to injection, stems were trimmed
 and flushed as described above for K_h measurements. The

basal end of the segments were then injected with a 10:1 silicone hardener mixture (Rhodorsil RTV 141; Bluestar Silicones, distributed by Skycon, Toronto, ON, Canada) that was mixed with a soluble fluorescent dye (1% w/w; Uvitex OB, Ciba Specialty Chemicals, Tarrytown, NY, USA dissolved in chloroform) at 50 kPa for 24 h. Stem segments were cured at room temperature for at least 72 h before sectioning. Stems were sectioned at several distances from the injection surface, and cross-sections were analysed to determine the percentage of vessels filled. The vessel length distribution was calculated based on the equations described in Wheeler *et al.* (2005), and this distribution was used to estimate the percentage of vessels open all of the way through segments and percentage of vessels open to the middle of segments from either cut end.

Statistical analysis

Plant material was carefully matched within each experiment, and vulnerability curves were compared within each experiment. We tested if dehydration and matched centrifuge vulnerability curves differed by comparing models that included both curves using nested models constrained to fit a single curve (Crawley 2007). Models were fitted to mean PLC of stems ($n = 1-3$) at each branch equilibrium or centrifuge xylem pressure. A reparameterised Weibull model (Ogle *et al.* 2009) was fitted to *Q. wislizeni* values using generalised nonlinear least squares, and incorporated a variance model to account for non-constant variance (Pinheiro & Bates 2000). Models were fitted with maximum likelihood, and comparison of nested models resulted in a likelihood ratio test statistic (LRT). The shape of the relationship between PLC and xylem pressure for *H. arbutifolia* was poorly fitted with the Weibull model, so a generalised additive model (GAM) was fitted (Wood 2006). This type of model does not assume a specific functional form for the shape of the relationship between response and explanatory variables.

We tested if centrifuge vulnerability curves constructed with 140 or 271-mm stems differed using two separate analyses. In both analyses, curves were corrected for cavitation fatigue (Hacke *et al.* 2001) by calculating the PLC using the K_h measured following the initial spin at 0.5 MPa in place of the maximum K_h . This minimised the effect of xylem vessels unlikely to be functional in intact plants. To compare P_{50} between vulnerability curves, we employed a common approach of fitting separate curves (GAM for *H. arbutifolia* and Weibull for other species) to data for each stem and then used the P_{50} estimates from fitted curves in an analysis of variance (ANOVA). We included species, stem length (140 mm or 271 mm) and their interaction in the model and tested for differences by stem length within species using *a priori* contrasts. A second approach that tested if the entire curves differed by stem length consisted of comparing models fitted to all stems that included two curves or were constrained to fit a single curve. This analysis was analogous to the dehydration and centrifuge curve comparison, but random effects were incorporated to account for the repeated measurements made on individual stems (Pinheiro & Bates 2000; Wood 2006).

The PLC response to repeated centrifugation was analysed as a mixed-model ANOVA. We included centrifugation (repeat number), vessel length (long or short) and centrifugation type

(repeated or single centrifugation) as fixed effects, and species and stem as random effects. Initial and final flushed K_s were similarly analysed by substituting flushed (initial or final) for centrifugation, and examined if initial and final K_s differed using *a priori* contrasts. For the repeated centrifugation experiment that tested the effect of foam pads in reservoirs, the PLC response to repeated centrifugation was analysed as a mixed-model ANOVA that included centrifugation (repeat number) and treatment (with foam or without foam) as fixed effects, and stem as a random effect. The first and the fourth centrifugation PLCs for each treatment were compared using *a priori* contrasts. Additional treatments tested for their effect on PLC induced by repeated centrifugation were modelled similarly. The analyses were performed using R version 2.8.1 (R Development Core Team 2011) including packages nlme (Pinheiro *et al.* 2011) and mgcv (Wood 2006).

RESULTS

Comparison of dehydration and centrifuge methods

Dehydration and centrifuge methods for inducing negative xylem pressure in stems were compared by constructing matched vulnerability curves for stems that were initially dehydrated and subsequently flushed of emboli and centrifuged. For both long-vesselled *Q. wislizeni* and short-vesselled *H. arbutifolia*, the shape of the centrifuge vulnerability curve was similar to the dehydration vulnerability (Fig. 2). The P_{50} for *Q. wislizeni* did not differ between dehydration and centrifuge curves (Table 2), but centrifugation produced modestly higher PLC than dehydration at a PLC >60% (Fig. 2a), resulting in differences between curves compared along their entire span ($LRT_2 = 31.85$, $P < 0.001$). Native PLC of *Q. wislizeni* stems sampled at the time of year when plants had high and low pre-dawn water potential closely matched the vulnerability curves (Fig. 2a). For *H. arbutifolia*, the dehydration and centrifuge vulnerability curves did not differ (Fig. 2b; $F_{1,8, 52.1} = 2.68$, $P = 0.083$), nor did P_{50} values (Table 2). Thus, in contrast to results of Cochard *et al.* (2010) (Fig. 1), the cavitation response of stems did not diverge widely depending on whether negative xylem pressure was generated by either centrifugation or air-drying large branches or natural dehydration in the field.

As a second independent measure of cavitation, we measured ultrasonic acoustic emissions emitted by xylem tissue during dehydration vulnerability curves. A high rate of acoustic emissions was observed early in the dehydration of *Q. wislizeni* branches as xylem pressure began to decrease, but thereafter emissions declined to a low rate, even as xylem pressure continued to drop (Fig. 3). This pattern of acoustic emissions corroborates the shape of the *Q. wislizeni* vulnerability curves, which predict extensive cavitation at the xylem pressures where the rate of acoustic emissions was high, but less additional cavitation at more negative xylem pressures where the acoustic emissions rate was low.

While constructing dehydration vulnerability curves, we found that xylem pressure potential of branchlets from the equilibrated branches measured with the pressure chamber agreed with stem psychrometer water potential measurements for *H. arbutifolia*, but were more negative at water potentials below ~ -4 MPa for *Q. wislizeni* (Figure S1).




Fig. 2. Percentage loss of hydraulic conductivity (PLC) as a function of xylem pressure induced by laboratory dehydration (closed circles) or subsequent centrifugation (open circles) of the same stems for long-vesselled *Quercus wislizeni* and shorter-vesselled *Heteromeles arbutifolia*. Points are means ($n = 1-3$) and lines are Weibull model fits for *Q. wislizeni* and generalised additive model fits for *H. arbutifolia*. Bars are native PLC of *Q. wislizeni* stems at shrub pre-dawn water potential measured during the summer dry season, 2009, and early spring wet season of 2010.

Table 2. The xylem pressure at 50% loss of hydraulic conductivity (P_{50}) of *Quercus wislizeni* and *Heteromeles arbutifolia* calculated from matched vulnerability curves constructed for stems initially dehydrated and subsequently flushed of emboli and centrifuged.

vulnerability curve	P_{50} (MPa)	
	<i>Quercus wislizeni</i>	<i>Heteromeles arbutifolia</i>
Dehydration	-1.35 (-1.71, -0.99)	-1.44 (-1.96, -0.99)
Centrifugation	-1.37 (-2.17, -0.59)	-2.09 (-5.22, -1.50)

Values shown are means and 95% confidence intervals in parentheses.




Fig. 3. The rate of acoustic emission and stem xylem water potential for a *Quercus wislizeni* branch during laboratory dehydration ($n = 3$ for stem water potential means; closed circles).

Comparison of centrifuge vulnerability curves of two stem lengths

To test if the proportion of open vessels affected vulnerability curves measured with the centrifuge method, we compared curves for stems of 140 and 271-mm lengths. For *H. arbutifolia*, shortening the length of stem segments from 271 to 140 mm only substantially changed the percentage of vessels open to the middle, but both categories of open vessels changed substantially for *Q. wislizeni* (Table 1). Changes in percentage of open vessels between stem lengths for *V. vinifera* were similar to *Q. wislizeni* (e.g. entirely open changed from 1.5% to 9.4%;

Jacobsen & Pratt 2012) and may also be similar for *Q. robur*, which has a similar maximum vessel length to *Q. wislizeni*. In spite of these changes, P_{50} did not differ by segment length for any of the four species examined (Table 3; $F_{1,50} < 2.28$, $P > 0.137$ for all segment length contrasts within species). Note that the P_{50} values in Tables 2 and 3 cannot be compared directly, because Table 3 values were calculated from vulnerability curves corrected for cavitation fatigue (Hacke *et al.* 2001), a correction not possible for dehydration vulnerability curves. In a separate test for differences along the entire span of vulnerability curves, the curves for the two stem lengths did not differ for *Q. wislizeni* or *H. arbutifolia* (Fig. 4; $LRT_2 = 3.36$, $P = 0.187$ and $LRT_3 = 0.02$, $P = 0.999$, respectively), but did for *Q. robur* and *V. vinifera* ($LRT_2 = 8.99$, $P = 0.011$, $LRT_3 = 7.48$, $P = 0.024$, respectively). The magnitude of the difference between vulnerability curves of the two stem lengths for *Q. robur* and *V. vinifera* were small compared to differences in vulnerability curves related to stem length observed by Cochard *et al.* (2010) for *Prunus persica* and *Q. robur* (e.g. *Q. robur*, Fig. 1).

Distribution of centrifugation-induced embolism

We compared the distribution of embolism along stems with the theoretical negative pressure profile induced by centrifugation for a long-vesselled species. The PLC of stem segments excised from the centre of a centrifuged stem was higher than the PLC of segments excised from the ends (Fig. 5). This distribution of PLC matches the expected effect of the lower negative pressure generated near the axis of rotation at the centre of the stem during centrifugation (Alder *et al.* 1997).

Repeated centrifugation tests

Repeated centrifugation of stems at the same angular velocity led to increases in PLC with each additional centrifugation for six chaparral shrub species, but the magnitude of these increases was not related to having long or short vessels (Figure S2). For all species, flushing following repeated centrifugation fully recovered the initial flushed maximum K_h . Centrifugal force maintained solution in contact with stem ends during centrifugation, but stem ends were exposed to air when the rotor was not spinning. We substantially reduced the PLC

Table 3. The xylem pressure at 50% loss of hydraulic conductivity (P_{50}) calculated from fatigue-corrected centrifuge vulnerability curves constructed with stem segments of two lengths for *Quercus wislizeni*, *Heteromeles arbutifolia*, *Quercus robur* and *Vitis vinifera* (for *Vitis vinifera* $n = 10$ for 140-mm stems and $n = 12$ for 271-mm stems, $n = 6$ for all other species).

stem length (mm)	P_{50} (MPa)			
	<i>Quercus wislizeni</i>	<i>Heteromeles arbutifolia</i>	<i>Quercus robur</i>	<i>Vitis vinifera</i>
140	-1.39 (-1.96, -0.82)	-6.78 (-7.98, -5.59)	-1.67 (-2.36, -0.97)	-1.22 (-1.48, -0.96)
271	-1.72 (-2.42, -1.01)	-7.00 (-8.23, -5.78)	-2.26 (-2.54, -1.97)	-1.45 (-1.73, -1.17)

Values shown are means and 95% confidence intervals in parentheses.




Fig. 4. Percentage loss of hydraulic conductivity as a function of xylem pressure induced by centrifugation of 140-mm (closed circles) and 271-mm (open circles) stem segments for long-vesselled *Quercus wislizeni* and shorter-vesselled *Heteromeles arbutifolia*. Points are means with SE ($n = 6$).

increases during repeated centrifugation by adding foam pads to the reservoirs to maintain solution contact with stem ends at all times (Figure S3). We also tried adding 10 min at 0.1 MPa to the end of centrifugation as an equilibration period and additional degassing with a membrane contactor, but these treatments had no effect on PLC increases in *Q. wislizeni* and *R. ovata*, respectively (data not shown).

DISCUSSION

Centrifuge vulnerability curves agree with dehydration curves

We found no evidence for an artefact associated with open vessels for vulnerability curves constructed using a centrifuge technique. These findings contrast with those of Cochard *et al.* (2010) and Choat *et al.* (2010). The vulnerability curves resulting from laboratory dehydration or centrifugation were similar for both a long- and a short-vesselled species, and other studies have found similar agreement between these methods (Alder *et al.* 1997; Jacobsen *et al.* 2007a; Jacobsen & Pratt 2012).

Vulnerability curves with a so-called 'r' shape, like the *Q. wislizeni* curve, characterised by large increases in PLC with the initial drops in xylem pressure have been called into question by Cochard *et al.* (2010). The centrifuge vulnerability curve shape we report for *Q. wislizeni* was supported by a matching laboratory dehydration curve, the pattern of acoustic

emissions of dehydrating branches and seasonal native embolism measurements. When employing the single-vessel air injection method (Melcher *et al.* 2003), which is a completely different methodology to that used here, Christman *et al.* (2012) found that another *Quercus* species had xylem with a similarly high proportion of highly vulnerable vessels. The 'r' shape has been previously observed for dehydration vulnerability curves of several other chaparral species (Jacobsen *et al.* 2007b), suggesting that multiple species have a high proportion of very vulnerable xylem vessels in this plant community. These highly vulnerable vessels may be functional and enhance carbon gain primarily during the wet season, or they may undergo daily cavitation and refilling cycles (Taneda & Sperry 2008; Christman *et al.* 2012), thereby maximising daily photosynthesis throughout the growing season. Alternatively, highly vulnerable vessels may be an efficient way to construct xylem having the required mechanical support.

The proportion of open vessels does not affect centrifuge vulnerability curves

We found no effect of changing the stem segment length and thereby the proportion of open vessels on the P_{50} of centrifuge vulnerability curves. The lack of a substantial stem length effect for long-vesselled species, a finding consistent with Sperry *et al.* (2012) and Jacobsen & Pratt (2012), does not support the




Fig. 5. Percentage loss of hydraulic conductivity of basal, central and distal 40-mm segments cut from 271-mm long *Rhus ovata* stems ($n = 3$) centrifuged to induce -2 MPa of pressure at the axis of rotation at the centre of the stem.

hypothesis that open vessels exhibit more vulnerability to cavitation during centrifugation. As described in Sperry *et al.* (2012), the increased vulnerability to cavitation of shorter stems for medium- to long-vesselled species reported by Cochard *et al.* (2010) may result from micro-bubbles being carried into the stem by the flow of solution induced as part of the unique Cavitron measurement technique. In contrast to Choat *et al.* (2010), we saw no evidence for an open-vessel artefact in centrifuge vulnerability curves of *V. vinifera*. A recent study suggests that *V. vinifera* branches used in dehydration vulnerability curves can develop vessel blockages, causing dehydration and centrifuge vulnerability curves to differ (Jacobsen & Pratt 2012).

Another result cited in Cochard *et al.* (2010) as demonstrating increased vulnerability of open vessels was the shift in Cavitron vulnerability curves following injection of air into the ends of stem segments. However, dramatically reducing hydraulic conductivity at the ends of stem segments with air injection (see Fig. 5 in Cochard *et al.* 2010) will shift vulnerability curves to appear more resistant to cavitation, regardless of which subset of vessels remain conductive. Centrifugation reduces hydraulic conductivity primarily near the middle of the segment length; therefore, a decrease in conductivity due to centrifugation results in a smaller proportional decrease in total stem conductivity when segment ends have been air-injected.

The pattern of cavitation is consistent with theoretical predictions

The differences between the distribution of embolism along the length of centrifuged stems measured in this study and that of Cochard *et al.* (2010) may partially explain why we do not observe a similar effect of stem length on centrifuge vulnerability curves. The distribution of PLC we measured for a long-vesselled species matches the theoretical expectation that embolism of a stem following centrifugation should be highest at the axis of rotation in the central part of the segment where the induced xylem pressure is most negative (Alder *et al.* 1997). Similar measurements conducted with the Cavitron found higher PLC at the basal stem end rather than at the

centre (Fig. 5 in Cochard *et al.* 2010). Cochard *et al.* (2010) suggest this distribution could result from vessel draining, but the match between our results and the predicted pattern of PLC suggest that vessel draining is an unlikely explanation. Sperry *et al.* (2012) describe the high PLC at the basal stem end as likely arising from micro-bubbles or debris carried into the stem by the flow of solution during Cavitron measurements. The centrifuge vulnerability curve technique we used does not induce flow during centrifugation and therefore would not be vulnerable to this artefact.

CONCLUSION

We conclude that the vulnerability of plants to cavitation can be accurately characterised with vulnerability curves constructed using a centrifuge technique, even for some long-vesselled species. For long- and short-vesselled species, the centrifuge technique used in this study did not produce the large artefacts reported in Cochard *et al.* (2010) for medium- and long-vesselled species using the Cavitron technique. Our results are consistent with other recent studies that have investigated the accuracy of centrifuge vulnerability curves (Jacobsen & Pratt 2012; Sperry *et al.* 2012).

ACKNOWLEDGEMENTS

We thank Samuel Del Rio and Evan MacKinnon for laboratory assistance. We thank John Sperry and Stephen Davis for valuable discussions relevant to this study, and Katie VinZant of the National Forest Service for help in obtaining research permits. Funding was provided by the National Science Foundation CAREER award (IOS-0845125) and the Andrew W. Mellon Foundation.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article:

Figure S1. Comparison of stem xylem water potential measured with a stem psychrometer and branchlet pressure potential measured with a pressure chamber ($n = 4-6$ for branchlet means) for large branches of *Quercus wislizeni* and *Heteromeles arbutifolia* equilibrated for at least 16 h.

Figure S2. The PLC of stems repeatedly centrifuged four times at 0.5 MPa for 5 min or once for 20 min and then flushed to remove embolism for three long-vesselled (*Quercus wislizeni*, *Rhus ovata*, *Fraxinus dipetala*) and three short-vesselled species (*Heteromeles arbutifolia*, *Ceanothus leucodermis*, *Ribes malvaceum*).

Figure S3. The PLC of *Rhus ovata* stems repeatedly centrifuged four times at 0.5 MPa for 5 min with standard reservoirs (closed circles) or with hydration reservoirs (open circles) that had foam pads added and then flushed to remove embolism. Points are means with SE ($n = 6$).

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

REFERENCES

- Alder N.N., Pockman W.T., Sperry J.S., Nuismer S. (1997) Use of centrifugal force in the study of xylem cavitation. *Journal of Experimental Botany*, **48**, 665–674.
- Brodribb T., Holbrook N.M. (2004) Stomatal protection against hydraulic failure: a comparison of coexisting ferns and angiosperms. *New Phytologist*, **162**, 663–670.
- Choat B., Drayton W.M., Brodersen C., Matthews M. A., Shackel K.A., Wada H., McElrone A.J. (2010) Measurement of vulnerability to water stress-induced cavitation in grapevine: a comparison of four techniques applied to a long-vesseled species. *Plant, Cell and Environment*, **33**, 1502–1512.
- Christman M.A., Sperry J.S., Smith D.D. (2012) Rare pits, large vessels, and extreme vulnerability to cavitation in a ring-porous tree species. *New Phytologist*, **193**, 713–720.
- Cochard H. (2002) A technique for measuring xylem hydraulic conductance under high negative pressures. *Plant, Cell and Environment*, **25**, 815–819.
- Cochard H., Damour G., Bodet C., Tharwat I., Poirier M., Améglio T. (2005) Evaluation of a new centrifuge technique for rapid generation of xylem vulnerability curves. *Physiologia Plantarum*, **124**, 410–418.
- Cochard H., Herbette S., Barigah T., Badel E., Ennajeh M., Vilagrosa A. (2010) Does sample length influence the shape of xylem embolism vulnerability curves? A test with the Cavitrone spinning technique. *Plant, Cell and Environment*, **33**, 1543–1552.
- Crawley M.J. (2007) *The R book*. John Wiley & Sons Inc., Hoboken, NJ, USA, 942 pp.
- Dixon H.H., Joly J. (1895) On the ascent of sap. *Philosophical Transactions of the Royal Society London, B Biological Sciences*, **186**, 563–576.
- Espino S., Schenk H.J. (2011) Mind the bubbles: achieving stable measurements of maximum hydraulic conductivity through woody plant samples. *Journal of Experimental Botany*, **62**, 1119–1132.
- Hacke U.G., Stiller V., Sperry J.S., Pittermann J., McCulloh K. (2001) Cavitation fatigue: embolism and refilling cycles can weaken the cavitation resistance of xylem. *Plant Physiology*, **125**, 779–786.
- Holbrook N.M., Burns M.J., Field C.B. (1995) Negative xylem pressures in plants: a test of the balancing pressure technique. *Science*, **270**, 1193–1194.
- Jackson G.E., Grace J. (1996) Field measurements of xylem cavitation: are acoustic emissions useful? *Journal of Experimental Botany*, **47**, 1643–1650.
- Jacobsen A.L., Pratt R.B. (2012) No evidence for an open vessel effect in centrifuge-based vulnerability curves of a long vesselled liana (*Vitis vinifera*). *New Phytologist*, **194**, 982–990.
- Jacobsen A.L., Pratt R.B., Davis S.D., Ewers F.W. (2007a) Cavitation resistance and seasonal hydraulics differ among three arid Californian plant communities. *Plant, Cell and Environment*, **30**, 1599–1609.
- Jacobsen A.L., Pratt R.B., Ewers F.W., Davis S.D. (2007b) Cavitation resistance among twenty-six chaparral species of southern California. *Ecological Monographs*, **77**, 99–115.
- Kursar T.A., Engelbrecht B.M.J., Burke A., Tyree M.T., El Omari B., Giraldo J.P. (2009) Tolerance to low leaf water status of tropical tree seedlings is related to drought performance and distribution. *Functional Ecology*, **23**, 93–102.
- Li Y., Sperry J.S., Taneda H., Bush S.E., Hacke U.G. (2008) Evaluation of centrifugal methods for measuring xylem cavitation in conifers, diffuse- and ring-porous angiosperms. *New Phytologist*, **177**, 558–568.
- Maherali H., Pockman W.T., Jackson R.B. (2004) Adaptive variation in the vulnerability of woody plants to xylem cavitation. *Ecology*, **85**, 2184–2199.
- Melcher P.J., Zwieniecki M.A., Holbrook N.M. (2003) Vulnerability of xylem vessels to cavitation in sugar maple. Scaling from individual vessels to whole branches. *Plant Physiology*, **131**, 1775–1780.
- Melcher P.J., Holbrook N.M., Burns M.J., Zwieniecki M.A., Cobb A.R., Brodribb T.J., Choat B., Sack L. (2012) Measurements of stem xylem hydraulic conductivity in the laboratory and field. *Methods in Ecology and Evolution*, **3**, 1111/1112. doi: 10.1111/j.2041-210X.2012.00204.x.
- Ogle K., Barber J.J., Willson C., Thompson B. (2009) Hierarchical statistical modeling of xylem vulnerability to cavitation. *New Phytologist*, **182**, 541–554.
- Pinheiro J.C., Bates D.M. (2000) *Mixed-effect models in S and S-PLUS*. Springer-Verlag, New York, NY, 528 pp.
- Pinheiro J.C., Bates D.M., DebRoy S., Sarkar D., the R Development Core Team (2011) nlme: Linear and nonlinear mixed effect models. R package version 3.1-101.
- Pockman W.T., Sperry J.S. (2000) Vulnerability to xylem cavitation and the distribution of Sonoran Desert vegetation. *American Journal of Botany*, **87**, 1287–1299.
- Pockman W.T., Sperry J.S., O'Leary J.W. (1995) Sustained and significant negative water pressure in xylem. *Nature*, **378**, 715–716.
- Pratt R.B., Jacobsen A.L., Golgotiu K.A., Sperry J.S., Ewers F.W., Davis S.D. (2007) Life history type and water stress tolerance in nine California chaparral species (Rhamnaceae). *Ecological Monographs*, **77**, 239–253.
- Pratt R.B., Jacobsen A.L., Mohla R., Ewers F.W., Davis S.D. (2008) Linkage between water stress tolerance and life history type in seedlings of nine chaparral species (Rhamnaceae). *Journal of Ecology*, **96**, 1252–1265.
- R Development Core Team (2011) *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Sperry J.S., Tyree M.T. (1988) Mechanism of water stress-induced xylem embolism. *Plant Physiology*, **88**, 581–587.
- Sperry J.S., Donnelly J.R., Tyree M.T. (1988a) A method for measuring hydraulic conductivity and embolism in xylem. *Plant, Cell and Environment*, **11**, 35–40.
- Sperry J.S., Tyree M.T., Donnelly J.R. (1988b) Vulnerability of xylem to embolism in a mangrove vs an inland species of Rhizophoraceae. *Physiologia Plantarum*, **74**, 276–283.
- Sperry J.S., Hacke U.G., Wheeler J.K. (2005) Comparative analysis of end wall resistivity in xylem conduits. *Plant, Cell and Environment*, **28**, 456–465.
- Sperry J.S., Christman M.A., Torres-Ruiz J.M., Taneda H., Smith D.D. (2012) Vulnerability curves by centrifugation: is there an open vessel artefact, and are 'r' shaped curves necessarily invalid? *Plant, Cell and Environment*, **35**, 601–610.
- Taneda H., Sperry J.S. (2008) A case-study of water transport in co-occurring ring- versus diffuse-porous trees: contrasts in water-status, conducting capacity, cavitation and vessel refilling. *Tree Physiology*, **28**, 1641–1651.
- Tyree M.T., Sperry J.S. (1989) Vulnerability of xylem to cavitation and embolism. *Annual Review of Plant Physiology and Plant Molecular Biology*, **40**, 19–38.
- Wheeler J.K., Sperry J.S., Hacke U.G., Hoang N. (2005) Inter-vessel pitting and cavitation in woody Rosaceae and other vesselled plants: a basis for safety versus efficiency trade-off in xylem transport. *Plant, Cell and Environment*, **28**, 800–812.
- Wood S.N. (2006) *Generalized additive models: an introduction with R*. Chapman and Hall/CRC Press, Boca Raton, FL, USA, 384 pp.

Author Query Form

Journal: PLB

Article: 678

Dear Author,

During the copy-editing of your paper, the following queries arose. Please respond to these by marking up your proofs with the necessary changes/additions. Please write your answers on the query sheet if there is insufficient space on the page proofs. Please write clearly and follow the conventions shown on the attached corrections sheet. If returning the proof by fax do not write too close to the paper's edge. Please remember that illegible mark-ups may delay publication.

Many thanks for your assistance.

Query reference	Query	Remarks
1	AUTHOR: Please provide the volume number, page first for reference Melcher et al. (2012).	
2	AUTHOR: Please provide missing details if any for ref. Pinheiro et al. (2011).	
3	AUTHOR: Since 2011, Research Papers will be subject to a page charge at a rate of GBP 100 per page above the 8-page limit (the first 8 pages are free of charge). Your article exceeds the page limit. Please go to the journal website, download the Page Charges Form, fill it in and send it to the Production Editor at plb@wiley.com indicating the paper number. (Please, click on the following link to download the form: http://www.blackwellpublishing.com/pdf/PLB_Page_Charge_Form.pdf).	

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ [Ⓢ]
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ [Ⓢ]
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↙
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Y or Y under character e.g. Y or Y
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	Y or Y and/or Y or Y
Insert double quotation marks	(As above)	Y or Y and/or Y or Y
Insert hyphen	(As above)	H
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	┌┐	┌┐
Close up	linking ○ characters	○
Insert or substitute space between characters or words	/ through character or ∧ where required	Y
Reduce space between characters or words		↑